

CONSTRUCTION OF NORMAL NUMBERS WITH RESPECT TO GENERALIZED LÜROTH SERIES FROM EQUIDISTRIBUTED SEQUENCES

MAX AEHLE AND MATTHIAS PAULSEN

ABSTRACT. Generalized Lüroth series generalize b -adic representations as well as Lüroth series. Almost all real numbers are normal, but it is not easy to construct one. In this paper, a new construction of normal numbers with respect to Generalized Lüroth Series (including those with an infinite digit set) is given. Our method concatenates the beginnings of the expansions of an arbitrary equidistributed sequence.

1. INTRODUCTION

In 1909, Borel proved that the b -adic representations of almost all real numbers are normal for any integer base b . However, only a few simple examples for normal numbers are known, like the famous Champernowne constant [Cha33]

$$0.1234567891011121314151617181920\dots$$

Other constants as π or e are conjectured to be normal. The concept of normality naturally extends to more general number expansions, including Generalized Lüroth Series [DK02, pp. 41–50]. For example, the numbers $1/e$ and $\frac{1}{2}\sqrt{3}$ are not normal with respect to the classical Lüroth expansion that was introduced in [Lü83].

In his Bachelor thesis, Boks [Bok09] proposed an algorithm that yields a normal number with respect to the Lüroth expansion, but he did not complete his proof of normality. Some years later, Madritsch and Mance [MM14] transferred Champernowne’s construction to any invariant probability measure. However, their method is rather complicated and does not reflect that digit sequences represent numbers. Vandehey [Van14] provided a much simpler construction, but for finite digit sets only, in particular, not for the original Lüroth expansion.

Like classical Lüroth expansions, continued fraction expansions are number representations with an infinite digit set. Adler, Keane and Smorodinsky [AKS81] showed that the concatenation of the continued fractions of

$$(1) \quad \frac{1}{2}, \frac{1}{3}, \frac{2}{3}, \frac{1}{4}, \frac{2}{4}, \frac{3}{4}, \frac{1}{5}, \frac{2}{5}, \frac{3}{5}, \frac{4}{5}, \dots$$

leads to a normal number.

This motivates our approach for the construction of normal numbers with respect to Generalized Lüroth Series (GLS) from equidistributed sequences. The idea is to look at the expansions of some equidistributed sequence with respect to a given

Date: September 28, 2015.

2010 Mathematics Subject Classification. Primary 11K16, Secondary 11A63.

Key words and phrases. Normal numbers, Generalized Lüroth Series, Equidistributed sequences, Numeration systems.

GLS. In general, these expansions are infinite. Hence, in order to concatenate them, we have to trim each digit sequence to a certain length.

The paper is organized as follows: In Sections 2 and 3 we recall the notions of equidistributed sequences and GLS respectively, then in Section 4 we describe our construction, finally, in Section 5 we prove that the constructed number is normal.

2. EQUIDISTRIBUTED SEQUENCES

Recall that a sequence (a_1, a_2, \dots) of numbers $a_j \in [0, 1]$ is called equidistributed if for any interval $I \subseteq [0, 1]$,

$$(2) \quad \lim_{n \rightarrow \infty} \frac{\#\{1 \leq j \leq n : a_j \in I\}}{n} = |I|.$$

Remark 1. If (a_1, a_2, \dots) is equidistributed, then (a_k, a_{k+1}, \dots) is equidistributed for any $k \in \mathbb{N} = \{1, 2, \dots\}$.

Many easy computable equidistributed sequences exist, e.g. the sequence (1) used by [AKS81] and the sequence $(n\beta \bmod 1)_{n \in \mathbb{N}}$ for any $\beta \in \mathbb{R} \setminus \mathbb{Q}$ [KN74].

It is convenient to introduce the concept of uniform equidistribution:

Definition 2. A sequence (a_1, a_2, \dots) of numbers $a_j \in [0, 1]$ is called *uniformly equidistributed* if the convergence in equation (2) is uniform in I , i.e. for all $\varepsilon > 0$ there is an $N \in \mathbb{N}$ such that for all $n \geq N$ and any interval $I \subseteq [0, 1]$,

$$\left| \frac{\#\{1 \leq j \leq n : a_j \in I\}}{n} - |I| \right| < \varepsilon.$$

As it turns out, these two terms are equivalent. This fact is commonly used without proving it. For clarity, we will sketch a simple proof here.

Lemma 3. *Every equidistributed sequence is uniformly equidistributed.*

Proof. Consider an equidistributed sequence (a_1, a_2, \dots) . Let $\varepsilon > 0$. Choose $k \in \mathbb{N}$ such that $\frac{1}{k} < \varepsilon/3$. Set $E_i = \left[\frac{i-1}{k}, \frac{i}{k}\right)$ for $i = 1, \dots, k-1$ and $E_k = \left[\frac{k-1}{k}, 1\right]$. For each E_i there is an $N_i \in \mathbb{N}$ with

$$\left| \frac{\#\{1 \leq j \leq n : a_j \in E_i\}}{n} - |E_i| \right| < \frac{\varepsilon}{3k}$$

for all $n \geq N_i$. Let $N = \max\{N_1, \dots, N_k\}$. For a given interval $I \subseteq [0, 1]$, consider the intervals E_s, \dots, E_t ($1 \leq s \leq t \leq k$) which have a non-empty intersection with I . Take an arbitrary integer $n \geq N$. We get

$$\begin{aligned} \frac{\#\{1 \leq j \leq n : a_j \in I\}}{n} - |I| &\leq \frac{\#\{1 \leq j \leq n : a_j \in \bigcup_{i=s}^t E_i\}}{n} - \left| \bigcup_{i=s+1}^{t-1} E_i \right| \\ &\leq \sum_{i=s}^t \left(\frac{\#\{1 \leq j \leq n : a_j \in E_i\}}{n} - |E_i| \right) + |E_s| + |E_t| \\ &\leq \sum_{i=1}^k \left| \frac{\#\{1 \leq j \leq n : a_j \in E_i\}}{n} - |E_i| \right| + \frac{2}{k} \\ &< k \cdot \frac{\varepsilon}{3k} + 2\varepsilon/3 = \varepsilon. \end{aligned}$$

Similarly, $-\varepsilon$ is a lower bound. □

We are interested in maps which preserve the equidistribution property of sequences.

Definition 4. A map $T: [0, 1] \rightarrow [0, 1]$ is called *equidistribution-preserving* if for every equidistributed sequence (a_1, a_2, \dots) the sequence (Ta_1, Ta_2, \dots) is equidistributed.

3. GENERALIZED LÜROTH SERIES

Based on [DK02, pp. 41–50], we precise our notion of Generalized Lüroth Series.

Definition 5. Let $\mathcal{D} \subseteq \mathbb{N}$ be a non-empty set (called “digit set”). Suppose that for each $d \in \mathcal{D}$ there is an interval $\mathcal{I}_d \subseteq [0, 1]$ and a map $T_d: \mathcal{I}_d \rightarrow [0, 1]$ with the following properties:

- (1) $\bigcup_{d \in \mathcal{D}} \mathcal{I}_d = [0, 1]$;
- (2) the intervals \mathcal{I}_d are pairwise disjoint;
- (3) if $|\mathcal{I}_d| > 0$, then T_d is either of the form

$$T_d(x) = \frac{1}{|\mathcal{I}_d|}(x - \inf \mathcal{I}_d) \quad \text{or} \quad T_d(x) = 1 - \frac{1}{|\mathcal{I}_d|}(x - \inf \mathcal{I}_d).$$

Define $T: [0, 1] \rightarrow [0, 1]$ by $T(x) = T_d(x)$ for $x \in \mathcal{I}_d$.

The tuple $\mathfrak{L} = (\mathcal{D}, \{\mathcal{I}_d : d \in \mathcal{D}\}, T)$ is called a *Generalized Lüroth Series (GLS)*. For $j \in \mathbb{N}_0$ we denote the j -th digit of $x \in [0, 1]$ with respect to \mathfrak{L} by $\mathfrak{L}_j(x)$, i. e. the $d \in \mathcal{D}$ such that $T^j x \in \mathcal{I}_d$. We write $\mathfrak{L}_j^\ell(x) = (\mathfrak{L}_j(x), \mathfrak{L}_{j+1}(x), \dots, \mathfrak{L}_{j+\ell-1}(x))$.

Indeed, the transformation of a GLS possesses the desired property.

Lemma 6. *Let $\mathfrak{L} = (\mathcal{D}, \{\mathcal{I}_d : d \in \mathcal{D}\}, T)$ be a GLS. Then T is equidistribution-preserving.*

Proof. Let (a_1, a_2, \dots) be a uniformly equidistributed sequence (by Lemma 3). Let $I \subseteq [0, 1]$ and $\varepsilon > 0$ be arbitrary. Because of properties (1) and (2) of Definition 5 we know $\sum_{d \in \mathcal{D}} |\mathcal{I}_d| = 1$, i. e. there are intervals $\mathcal{I}_{d_1}, \dots, \mathcal{I}_{d_k}$ with

$$(3) \quad \sum_{i=1}^k |\mathcal{I}_{d_i}| > 1 - \varepsilon/2.$$

The set $[0, 1] \setminus \bigcup_{i=1}^k \mathcal{I}_{d_i}$ is a union of pairwise disjoint intervals $H_0, \dots, H_{k'}$ with $k' \leq k$. Let $J_i = (T^{-1}I) \cap \mathcal{I}_{d_i}$ for $i = 1, \dots, k$. Definition 5 implies that J_1, \dots, J_k are intervals. It is a well known fact that T preserves the Lebesgue measure. Hence

$$(4) \quad \sum_{i=1}^k |J_i| \leq |I| \leq \sum_{i=1}^k |J_i| + \sum_{i=0}^{k'} |H_i|.$$

Using Definition 2, there exists an integer $N \in \mathbb{N}$ such that for all $n \geq N$ we have

$$(5) \quad \left| \frac{\#\{1 \leq j \leq n : a_j \in X\}}{n} - |X| \right| < \frac{\varepsilon}{4k+2},$$

whereas $X \in \{J_1, \dots, J_k, H_0, \dots, H_{k'}\}$. For the upper bound we get for every $n \geq N$

$$\begin{aligned}
\frac{\#\{1 \leq j \leq n : Ta_j \in I\}}{n} - |I| &\leq \sum_{i=1}^k \frac{\#\{1 \leq j \leq n : a_j \in J_i\}}{n} \\
&\quad + \sum_{i=0}^{k'} \frac{\#\{1 \leq j \leq n : a_j \in H_i\}}{n} - |I| \\
&\stackrel{(4)}{\leq} \sum_{i=1}^k \left(\frac{\#\{1 \leq j \leq n : a_j \in J_i\}}{n} - |J_i| \right) + \sum_{i=0}^{k'} |H_i| \\
&\quad + \sum_{i=0}^{k'} \left(\frac{\#\{1 \leq j \leq n : a_j \in H_i\}}{n} - |H_i| \right) \\
&\stackrel{(3),(5)}{<} k \cdot \frac{\varepsilon}{4k+2} + \varepsilon/2 + (k'+1) \cdot \frac{\varepsilon}{4k+2} \leq \varepsilon.
\end{aligned}$$

For the lower bound we similarly obtain

$$\begin{aligned}
|I| - \frac{\#\{1 \leq j \leq n : Ta_j \in I\}}{n} &\leq |I| - \sum_{i=1}^k \frac{\#\{1 \leq j \leq n : a_j \in J_i\}}{n} \\
&\stackrel{(4)}{\leq} \sum_{i=0}^{k'} |H_i| + \sum_{i=1}^k \left(|J_i| - \frac{\#\{1 \leq j \leq n : a_j \in J_i\}}{n} \right) \\
&\stackrel{(3),(5)}{<} \varepsilon/2 + k \cdot \frac{\varepsilon}{4k+2} < \varepsilon. \quad \square
\end{aligned}$$

With respect to a GLS \mathfrak{L} , a number $x \in [0, 1]$ is normal if for any block $b = (b_1, \dots, b_r) \in \mathcal{D}^r$, we have

$$(6) \quad \lim_{n \rightarrow \infty} \frac{\#\{0 \leq j \leq n-1 : \mathfrak{L}_j^r(x) = b\}}{n} = \prod_{i=1}^r |\mathcal{I}_{b_i}|.$$

4. CONSTRUCTION

Fix a GLS $\mathfrak{L} = (\mathcal{D}, \{\mathcal{I}_d : d \in \mathcal{D}\}, T)$ and an equidistributed sequence (a_1, a_2, \dots) . We recursively define a strictly increasing sequence (c_0, c_1, c_2, \dots) of indices as follows. Let $c_0 = 0$. Assume c_0, \dots, c_ℓ are already chosen for some $\ell \in \mathbb{N}_0$. For $i = 0, \dots, \ell$ the sequences $(T^i a_{c_i+1}, T^i a_{c_i+2}, \dots)$ are uniformly equidistributed by Remark 1, Lemma 3 and Lemma 6. Therefore we can choose a $c_{\ell+1} > c_\ell$ such that for all integers $n \geq c_{\ell+1}$ and $0 \leq i \leq \ell$ and any interval $I \subseteq [0, 1]$ we have

$$(7) \quad \left| \frac{\#\{c_i+1 \leq j \leq n : T^i a_j \in I\}}{n - c_i} - |I| \right| < \frac{1}{\ell+1}.$$

Finally, we define $z \in [0, 1]$ such that its expansion with respect to \mathfrak{L} is the concatenation of $\mathfrak{L}_0^{l(1)}(a_1)$, $\mathfrak{L}_0^{l(2)}(a_2)$, $\mathfrak{L}_0^{l(3)}(a_3)$, etc., whereas $l(j)$ is chosen such that $c_{l(j)-1} + 1 \leq j \leq c_{l(j)}$. In particular, we have $l(j) \leq j$ as well as $l(j) \rightarrow \infty$ for $j \rightarrow \infty$.




FIGURE 1. Visualization of the construction of z . The m -th digit of z in the i -th row ($i = 0, 1, \dots$) and j -th column ($j = 1, 2, \dots$) is $\mathfrak{L}_m(z) = \mathfrak{L}_i(a_j)$.

5. PROOF OF NORMALITY

Theorem 7. *The number z is normal with respect to \mathfrak{L} .*

Proof. Let $b = (b_1, \dots, b_r) \in \mathcal{D}^r$ be a block of digits. Define $I \subseteq [0, 1]$ as the set of all $x \in [0, 1]$ which fulfill $\mathfrak{L}_0^r(x) = b$. It follows that I is an interval of length $\prod_{i=1}^r |\mathcal{I}_{b_i}|$. We want to verify equation (6). Fix an arbitrary $n \in \mathbb{N}$. Assume that the $(n-1)$ -th digit of z (i.e. $\mathfrak{L}_{n-1}(z)$) occurs within $\mathfrak{L}_0^{l(k)}(a_k)$ for some $k \in \mathbb{N}$. Obviously, the number of digit positions $m \in \{0, \dots, n-1\}$ which belong to an $\mathfrak{L}_i(a_j)$ for a certain $i = 0, \dots, l(k)-1$ is bounded above by $k - c_i$, thus

$$(8) \quad n \leq \sum_{i=0}^{l(k)-1} (k - c_i) \leq \sum_{i=0}^{l(k)-2} (k - c_i) + k.$$

Similarly, it is bounded below by $k - 1 - c_i$, hence

$$(9) \quad n \geq \sum_{i=0}^{l(k)-1} (k - 1 - c_i) \geq \sum_{i=0}^{l(k)-2} (k - c_i - 1).$$

Counting the number of occurrences of b in each column separately (see Figure 1), we obtain

$$\begin{aligned} & \frac{\#\{0 \leq j \leq n-1 : \mathfrak{L}_j^r(z) = b\}}{n} - |I| \\ & \leq \sum_{j=1}^k \frac{\#\{0 \leq i \leq l(j) - r - 1 : \mathfrak{L}_i^r(a_j) = b\} + r}{n} - |I| \\ & \leq \sum_{j=1}^k \frac{\#\{0 \leq i \leq l(j) - 2 : T^i a_j \in I\}}{n} - |I| + \frac{k}{n} \cdot r \\ & = \sum_{i=0}^{l(k)-2} \frac{\#\{c_i + 1 \leq j \leq k : T^i a_j \in I\}}{n} - |I| + \frac{k}{n} \cdot r \\ & \stackrel{(9)}{\leq} \sum_{i=0}^{l(k)-2} \frac{k - c_i}{n} \left(\frac{\#\{c_i + 1 \leq j \leq k : T^i a_j \in I\}}{k - c_i} - |I| \right) + \frac{l(k) - 1}{n} \cdot |I| + \frac{k}{n} \cdot r \\ & \stackrel{(7)}{\leq} \sum_{i=0}^{l(k)-2} \frac{k - c_i}{n} \cdot \frac{1}{l(k) - 1} + \frac{l(k) - 1}{n} \cdot |I| + \frac{k}{n} \cdot r \end{aligned}$$

$$\stackrel{(9)}{\leq} \frac{n + l(k) - 1}{n(l(k) - 1)} + \frac{l(k) - 1}{n} \cdot |I| + \frac{k}{n} \cdot r \leq \frac{1}{l(k) - 1} + \frac{1}{n} + \frac{k}{n} \cdot (|I| + r).$$

In a similar manner, we establish a lower bound:

$$\begin{aligned} & |I| - \frac{\#\{0 \leq j \leq n-1 : \mathfrak{L}_j^r(z) = b\}}{n} \\ & \leq |I| - \sum_{i=0}^{l(k)-2} \frac{\#\{c_i + 1 \leq j \leq k-1 : T^i a_j \in I\}}{n} + \frac{k-1}{n} \cdot r \\ & \stackrel{(8)}{\leq} \sum_{i=0}^{l(k)-2} \frac{k - c_i - 1}{n} \left(|I| - \frac{\#\{c_i + 1 \leq j \leq k-1 : T^i a_j \in I\}}{k - c_i - 1} \right) \\ & \quad + \frac{l(k) - 1 + k}{n} \cdot |I| + \frac{k-1}{n} \cdot r \\ & \stackrel{(7),(9)}{\leq} \frac{1}{l(k) - 1} + \frac{l(k) - 1 + k}{n} \cdot |I| + \frac{k-1}{n} \cdot r \leq \frac{1}{l(k) - 1} + \frac{k}{n} \cdot (2|I| + r). \end{aligned}$$

For $n \rightarrow \infty$, we have $k \rightarrow \infty$ and thus $l(k) \rightarrow \infty$. Using the Cauchy limit theorem, we conclude

$$\frac{n}{k} \geq \frac{0 + l(1) + l(2) + \dots + l(k-1)}{k} \rightarrow \infty.$$

It follows that

$$\lim_{n \rightarrow \infty} \left| \frac{\#\{0 \leq j \leq n-1 : \mathfrak{L}_j^r(z) = b\}}{n} - |I| \right| = 0.$$

This completes the proof. \square

6. OPEN PROBLEMS

It would be more straightforward and closer to [AKS81], if the trimming of the expansions could be omitted by using equidistributed sequences of rational numbers with a finite expansion. Lüroth himself proved that rational numbers always have a finite or periodic Lüroth expansion [Lü83], but he did not find a way to distinguish between these two possibilities. Some algebraic manipulation shows that fractions of the form $\frac{a}{2^k}$ (i. e. the dyadic rationals) have a finite Lüroth expansion. We conjecture that this also applies to fractions of the form $\frac{a}{3^k}$. It would be very interesting to get a full understanding of rationals with a finite Lüroth expansion.

ACKNOWLEDGMENT

This paper was written during an internship at the Max Planck Institute for Mathematics in Bonn. We would like to thank the MPIM for offering us this unique opportunity to conduct active mathematical research. We are very grateful to our adviser, Izabela Petrykiewicz, for her helpful suggestions and support.

REFERENCES

- [AKS81] Roy Adler, Michael Keane, and Meir Smorodinsky, *A Construction of a Normal Number for the Continued Fraction Transformation*, *Journal of Number Theory* **13** (1981), 95–105.
- [Bok09] Arnout Boks, *Towards the construction of a normal Lüroth series*, Bachelor thesis, Technische Universiteit Delft, August 2009.

- [Cha33] David Gawen Champernowne, *The construction of decimals normal in the scale of ten*, Journal of the London Mathematical Society **8** (1933), 254–260.
- [DK02] Karma Dajani and Cor Kraaikamp, *Ergodic Theory of Numbers*, The Carus Mathematical Monographs, no. 29, The Mathematical Association of America, 2002.
- [KN74] Lauwerens Kuipers and Harald Niederreiter, *Uniform Distribution of Sequences*, Wiley, 1974.
- [Lü83] Jacob Lüroth, *Über eine eindeutige Entwicklung von Zahlen in eine unendliche Reihe*, Mathematische Annalen **21** (1883), 411–423.
- [MM14] Manfred Gerhard Madritsch and Bill Mance, *Construction of μ -normal Sequences*, arXiv:1206.4950v3 (2014).
- [NZ51] Ivan Niven and Herbert Samuel Zuckerman, *On the Definition of Normal Numbers*, Pacific Journal of Mathematics **1** (1951), 103–109.
- [Van14] Joseph Vandehey, *A Simpler Normal Number Construction for Simple Lüroth Series*, Journal of Integer Sequences **17** (2014).
- [Ša68] Tibor Šalát, *Zur metrischen Theorie der Lürothschen Entwicklungen der reellen Zahlen*, Czechoslovak Mathematical Journal **18** (1968), no. 93, 489–522.

MAX AEHLE, MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, D-53111 BONN
Current address: Faculty of Mathematics, Technische Universität Kaiserslautern, P. O. Box 3049,
D-67653 Kaiserslautern
E-mail address: aehle@rhrk.uni-kl.de

MATTHIAS PAULSEN, MAX PLANCK INSTITUTE FOR MATHEMATICS, VIVATSGASSE 7, D-53111 BONN
Current address: Department Mathematisches Institut, Universität München, Theresienstr. 39,
D-80333 München
E-mail address: paulsen@math.lmu.de